

THE EFFECTS OF CHEMICAL FERTILIZERS AND PESTICIDES ON THE ENVIRONMENT AND HUMAN HEALTH

Smt. Veena

Lecturer in Chimestry

S.N.D.B. Govt. College, Nohar

ABSTRACT

A surplus of agriculture is a vital factor that contributes to the development of a country's economy. It is also necessary to meet the increasing population's needs. The Green Revolution was created to meet these needs. The Green Revolution was able to help developing countries overcome their chronic food scarcity by increasing their agricultural production. This was done through the use of high-yielding seeds, the modification of farm equipment, and the use of chemical fertilizers. Although the use of certain agricultural practices allowed for the growth of food grains, they have also led to the development of harmful chemicals. This article talks about the effects of pesticides and chemical fertilizers on human health and the environment.

Keywords: Green revolution, chemical fertilizers, pesticides, environment, human health.

INTRODUCTION

The goal of agriculture revolves around meeting the present needs of the society while producing surplus supplies for export. To achieve this, the use of chemical inputs has increased. One of these is pesticides, which are used to kill pests. A chemical or biological agent that kills or deters harmful pests is known as a pesticide. It is commonly used to control or eliminate various agricultural pests that can severely affect crops and livestock. The use of pesticides has been beneficial for the farmers as it increases their agricultural yield. In addition, the Green Revolution has led to an increase in the number of pesticides being used in Indian agriculture, which helps the country fight against food crises. Although the use of pesticides can help prevent harmful pests from attacking crops, it can also cause long-term damage to the human and environmental health. India is currently the largest producer of these chemicals in Asia. Despite the lower consumption rate of pesticides in the country, the residue of these chemicals still remains a major issue.

Due to the increasing concerns about the hazards of chemicals used in agriculture, the government has started to consider the use of safer alternatives. This can be done through the development of better methods to reduce the use of pesticides. If the chemicals are used in the appropriate quantities and only when necessary, then the risks of exposure can be minimized.

Even though they are low in concentration, the use of pesticides has a serious effect on the environment. (Agrawal et al. 2010).

The data for the last two decades regarding pesticide exposure and human health revealed that several pesticides cause neuronal disorder and degenerative diseases, some effect fetal growth and cause congenital anomalies and other are carcinogenic for human (Asghar et al. 2016). Over the past three decades, the indiscriminate use and improper handling of pesticides in agriculture have caused serious human health problems in many developing countries (Dasgupta et al. 2007).

Effects of Chemical Fertilizers and Pesticides on Human Health

Bhandari (2014) in his study on the effects of agrochemicals on the environment in Nepal, Bhandari noted that they are considered as powerful tools that can be used to improve the productivity of agriculture in developing countries. However, it has been revealed that they are also causing various health issues and can lead to the development of cancer.

It has been administered that farmers do not use the safety masks, gloves and other protective gears during the spraying of pesticides which results into the access of pesticides in the blood stream through inhalation and dermal exposure which can adversely affect their eyes, skin and the respiratory system. The study shows relationship between the extent of pesticide used and signs and symptoms of illnesses due to exposure among spray farmers of Bhopal, Madhya Pradesh India, who sprayed pesticides by themselves and therefore were directly exposed to pesticides were assessed. The 18 months exposed spray farmers reported maximum acute signs and symptoms like burning/stinging of eyes (18.42%), blurred vision (23.68%), skin redness/itching (50%), excessive sweating/shortness of breath (34.2%), dry sore throat (21.05%) and burning of nose (28.9%). The signs and symptoms were found to be duration dependent among the sprayers. It is concluded that there is need for creating more awareness among the farm sprayers and authorities in implementing and ensuring the use of protective gear while handling pesticides (Choudhary 2014).

When fertilizers and pesticides are used in farmlands, they are transmitted directly or indirectly into the corns and vegetable that affects the human health. Moreover, as pesticides are applied over the vegetable which are directly entered into human or livestock bodies. Excessive use of fertilizers may pollute the underground water with nitrate and it is so much hazardous to humans or livestock. Nitrate concentrated water can immobilize some of hemoglobin in blood. Organophosphate pesticides have increased in application, because they are both less persistent and harmful for environment than organochlorin pesticides. But, they are associated with acute health problems, such as abdominal pain, dizziness, headaches, nausea, vomiting, as well as skin and eye problems. There have been many studies intending to establish cancer – pesticides association. Organophosphate pesticides used in the vegetables gradually get deposit into human body and has a link with cancer (Miah et al. 2014).

Wimalawansa and Wimalawansa (2014) assessed the impact of changing agricultural practices on human health in Srilanka and concluded that detrimental agricultural habits, including the excessive and indiscriminate use of toxic agrochemicals, allowing continued environmental contamination and contamination of the human food chain. Contamination of soil and water with toxic agrochemicals (e.g., phosphate fertilizer contaminated with heavy metals, pesticides and herbicides etc.) are a particular concern. These pollutants in water generally are in small quantities, and thus, cannot be seen or tasted. Therefore, their harmful effects do not manifest in humans for several years but led to the escalation of deadly disease like chronic kidney disease.

Green Revolution makes India self-dependent in term of food grains but indiscriminate use of synthetic fertilizers and pesticides contaminated our food and environment. Punjab, an agricultural state of the Indian Republic known as the grain bowl of the country is facing serious problems. Nutrient imbalance in the soil and surface water

contamination, pesticide residues in food and bovine milk and increasing cancer rate in farmers are some of the example of Green Revolution (Rahman and Debnath 2015).

The study shows that DDT was the most popular and effective pesticide to help people combat unwanted organisms and gain dramatically improvement in agriculture. However, since a number of adverse effects of this insecticide were reported, usage of DDT was banned international wide. Despite the severe restriction, DDT is still illegally used in many areas, especially in developing nations. Negative impacts of DDT on the human health were acknowledged and disseminated widely to warn population and prevent unexpected situations occur. Nevertheless, although DDT was not used in recent time, it still impacts on human health due to long residual efficacy and accumulation through food chain. In term of human health, DDT is the cause of many kinds of cancer, acute and persistent injury to the nervous system, lung damage, injury to the reproductive organs, dysfunction of the immune and endocrine systems, birth defects (Thuy, 2015).

The use of pesticides was introduced in India during the mid-sixties as a part of green revolution and malaria prevention programs. While pesticides turned useful for pest control they were at the same time responsible for human health injuries. Today these chemicals in particular those which accumulate in food chain, impose several human health hazards. Intake of food containing pesticide residues is documented to result in highest exposure, about 103 –105 times higher than that arising from contaminated drinking water or air. Pesticides have been reported to cause several adverse health effects which depend on the extent and duration of exposure. Health effects of pesticides range from mild allergies, rashes, breathing difficulties, neurotoxicity and reproductive abnormalities to deadly chronic diseases like cancer. This challenge to food safety may be addressed by preventive strategies which include the use of alternative sustainable agricultural practices or mitigating strategies which are based on reducing pesticide exposure from food and water by different processing techniques (Tomer et al. 2015).

Effects of chemical fertilizers and pesticides on environment

Soil, the basic need of farming may happen to pollute by the accumulation of various heavy metals, through emissions by industries, mining process, disposal of high metal wastes, gasoline, application of fertilizers, sewage sludge, pesticides, wastewater irrigation, coal combustion residues, etc. Historically, a large amount of chemicals is annually applied at the agricultural soils as fertilizers and pesticides. Such applications may result in the increase level of heavy metals, particularly Cd, Pb, and As in the soil (Atafar et al. 2010). Usage of pesticides, insecticides and other various chemicals in agriculture is very easy, quick and inexpensive solution for controlling weeds and insect pests. However, use of chemicals comes with a significant cost. They have contaminated almost every part of our environment and their residues are found in soil, water, land and air.

Kumar et al. (2013) concluded that pesticides are often considered a quick, easy and inexpensive solution for controlling weeds and insect pests in urban landscapes. Pesticides have contaminated almost every component of our environment. Pesticide residues are found in soil and air, and in surface and ground water across the nation, and urban pesticide uses contribute to the problem. Pesticide contamination poses significant risks to the environment and non-target organisms ranging from beneficial soil microorganisms, to insects, plants, fish, and birds. Contrary to common misconceptions, even herbicides can cause harm to the environment.

The study performed in the surface water of Sharda river region in Lakhimpurkheeri, Uttar Pradesh-India reports the concentration levels and distribution patterns of the 21 organochlorine pesticide residues in Solid Phase Extraction (SPE) is used for the extraction of organochlorine pesticide residues in water sample. The most commonly encountered Organochlorine pesticides in surface water were dieldrin, heptachlor epoxide, isomers of hexachlorocyclohexane and DDT. In some cases the concentrations detected were higher than the quantitative target levels set by the European Union, especially for γ -hexachlorocyclohexane & pp'-DDT. The concentration levels found are lower than the EU maximum acceptable concentration of 0.10 $\mu\text{g l}^{-1}$ for all compounds examined, except for δ -HCH in seven samples (0.2772, 0.1950, 0.2210, 0.2045, 0.1994, 0.1523, 0.1390 $\mu\text{g l}^{-1}$) and four samples (0.1877, 0.2365, 0.1478, 0.1269 $\mu\text{g l}^{-1}$) of pp'-DDT during 2008-2010.

The occurrence of these compounds in Sharda river region surface waters can be attributed to intense agricultural activity as well as to transboundary pollution (Maurya and Kumar 2013).

Pesticides have contaminated almost every part of our environment and pesticide residues are found in soil, air and in surface and groundwater. Pesticide contamination poses significant risks to the environment and non-target organisms ranging from beneficial soil microorganisms to insects, plants, fish, and birds. Recent studies have indicated that our environment is chronically polluted by pesticides and levels of biocidal contamination have increased tremendously. The environmental deterioration due to pesticides is endangering the situation of future (Sitaramaraju et al. 2014).

Soil analysis is carried out in the villages of Loni, Adgaon, Chinchpur, Sadatpur, Gogalgaon, falling in Taluka of Rahata & Sangamneer. Chemical fertilizers and pesticides are continuously being applied to agricultural fields for past many years boosting the agricultural yield and increasing chemical fertilizer consumption. The requirement of agricultural product (sugarcane, bajra, vegetables, food & fodder for animals) to cater to the domestic and industrial needs have made farmers to use fertilizer and pesticide beyond the prescribed dosage. The soil pH varied between from 7.46 to 8.9 and soil was found to be moderate alkaline. The soil is found to be free from salt accumulation. Organic carbon in soil was found to vary from moderate to very low indicating the need for improving the soil fertility. Available Nitrogen was found to be low in about 80% of soil sample, indicating lack of nitrogen status and the need for adequate application of nitrogenous fertilizer. 50% of the sample tested revealed very low Available Phosphorous content, remaining with moderate to low content. More than 80% of soil sample showed high value of Available Potassium as high as 963.2kg/ha, remaining have moderate to low values. Micronutrients, Zn, Cu, had moderate to low value and Fe showed very low value, but about 48% of sample had Mn higher value. This nature of micronutrients may probably due to the moderate alkali value in the soil Groundwater studies in the region have shown growing alkalinity, Nitrates and other fertilizer and pesticide residue in the water. The growing soil contamination can lead to further deterioration of ground water quality, which needs to be controlled through controlled application of water and application biofertilizer, organic manure (Natraj and Katyal 2014).

A research has been done in (Ardabil- Iran) Moghan's irrigation and drainage network in order to evaluate long term effects of pesticides and chemical fertilizers usage on soil properties and heavy metals accumulation. The results showed that soil physical characteristics such as bulk density were changed in long-term and it was increased compared to control soil. The heavy metals accumulations in soil were highly affected and the

concentration of some metals such as cadmium has reached a limit beyond the standard for agricultural purposes. The results also showed that fortunately the concentration of other metals is not beyond the standard. In this context, given the state of the soil and gained results, considering pesticides and fertilizers management is essential and it requires planning to reduce or replace pesticides and fertilizers usage in this region (Yargholi and Azarneshan 2014).

CONCLUSION

It has been concluded that many farmers do not follow proper safety precautions when it comes to the use of pesticides. This has led to the development of several human health issues and the pollution of our air, water, and land. Since a large portion of the population depends on agriculture for their livelihood, the use of pesticides is widely used to increase the production of food. There is a need to take adequate measures to protect the environment and human life from the harmful effects of pesticides. One of the most important steps that people can take to ensure that they are contributing to the environment is by adopting an organic farming system. This method of farming is an environmentally friendly way of doing things. By practicing organic agriculture, we are able to move back to our ancestors' course and contribute to the sustainability of the planet. This system of production and management supports the environment, human health, and food security. Despite the government's efforts to encourage people and farmers to adopt organic practices, it has not been able to bridge the gap between the demand for organic products and the supply of them in the market. It is important that the farmers are taught about the various aspects of organic farming to ensure that they are able to achieve their goals and protect the environment. Due to the increasing number of organic food stores in India, it has been observed that the demand for organic food is growing.

REFERENCES

- Abhilash, P.C. and Singh, N. 2008. Pesticide use and application: An Indian scenario. *Journal of Hazardous Materials*, 165: 1-12.
- Agrawal, A., Pandey, R.S. and Sharma, B. 2010. Water pollution with special reference to pesticide contamination in India. *Journal of Water Resource Protection*. 2(5): 432–448.
- Atafar, Z., Mesdaghinia, A., Nouri, J., Homae, M., Yunesian, M., Ahmadimoghaddam, M. and Mahvi, A.H. 2010. Effect of fertilizer application on soil heavy metal concentration. *Environment Monit Assess*, 160(1-4): 83-89.
- Choudhary, A., Ali, A.S. and Ali, S.A. 2014. Adverse Health Effects of Organophosphate Pesticides among Occupationally Exposed Farm Sprayers: A Case Study of Bhopal Madhya Pradesh, India. *Asian Journal of Biomedical and Pharmaceutical Sciences*, 4(35): 29-34
- Dasgupta, S., Meisner, C., Wheeler, D., Xuyen, K. and Lam, N.T. 2007. Pesticide Poisoning of farm workers- implications of blood test result from Vietnam. *International Journal of Hygiene Environment Health*, 210: 121-132.
- Dubey, R.K. 2013. Organic farming beneficial to biodiversity conservation, rural livelihood and nutritional security. *Indian Journal of Applied Research*, 3: 18-21.

- Kumar, S., Sharma, A.K., Rawat, S.S., Jain, D.K. and Ghosh, S. 2013. Use of pesticides in agriculture and livestock animals and its impact on environment of India. *Asian Journal of Environmental Science*. 8(1): 51-57.
- Maurya, A.K. and Kumar, A. 2013. Organochlorine Pesticides in the Surface Waters from Sharda River Region, Uttar Pradesh-India. *Advances in Space Research & Earth Exploration*, 1(1): 1-7.
- Miah, S.J., Hoque, A., Paul, A. and Rahman, A. 2014. Unsafe Use of Pesticide and Its Impact on Health of Farmers: A Case Study in Burichong Upazila, Bangladesh. *Journal of Environmental Science, Toxicology and Food Technology*, 8(1): 57-67.
- Natraj, V.M. and Katyaj, D. 2014. Study of Fertilizer Effect on soil status in and around Loni, Maharashtra, India. *Applied Sciences, Engineering and Technology*, 13: 188-192.
- Rahman, K.M. and Debnath, S.C. 2015. Agrochemical use, environmental and health hazards in Bangladesh. *International Research Journal of Interdisciplinary & Multidisciplinary Studies*, 1: 75-79.
- Sitaramaraju, S., Prasad, N.V.V.S.D., Chenga Reddy, V and Narayana, E. 2014. Impact of Pesticides Used for Crop Production on the Environment. *Journal of Chemical and Pharmaceutical Sciences*, 3: 75-79.
- Thuy, T.T. 2015. Effects of DDT on Environment and Human Health. *Journal of Education and Social Sciences*, 2: 108-114
- Tomer, V., Sangha, J.K. and Ramya, H.G. 2015. Pesticide: An Appraisal on Human Health Implications. *Proceedings of the National Academy of Sciences, India Section B: Biological Sciences*, 85(2): 451-463.
- Wimalawansa, S.A. and Wimalawansa, S.J. 2014. Impact of changing agricultural practices on human health: chronic kidney disease of multi-factorial origin in Sri Lanka. *Wudpecker Journal of Agricultural Research*, 3: 110-124.
- Yargholi, B. and Azarneshan, S. 2014. Long-term effects of pesticides and chemical fertilizers usage on some soil properties and accumulation of heavy metals in the soil (case study of Moghan plain's (Iran) irrigation and drainage network). *International Journal of Agriculture and Crop Sciences*, 7(8): 518-523.